



Excellence in CME
• April 5-9 •



Advertisement

Annals of Internal Medicine

Established in 1927 by the American College of Physicians

User Name

Password

LOG-IN

HOME | CURRENT ISSUE | PAST ISSUES | CME | AUDIO | ALERTS | COLLECTIONS | SUBSCRIBE | SUBMIT

Go

Advanced Search »

Follow Annals:    

Clinical Guidelines

Diagnosis and Treatment of Low Back Pain: A Joint Clinical Practice Guideline from the American College of Physicians and the American Pain Society

Roger Chou, MD; Amir Qaseem, MD, PhD, MHA; Vincenza Snow, MD; Donald Casey, MD, MPH, MBA; J. Thomas Cross, Jr, MD, MPH; Paul Shekelle, MD, PhD; Douglas K. Owens, MD, MS; and for the Clinical Efficacy Assessment Subcommittee of the American College of Physicians and the American College of Physicians/American Pain Society Low Back Pain Guidelines Panel*

Author Affiliations

Abstract

Recommendation 1: Clinicians should conduct a focused history and physical examination to help place patients with low back pain into 1 of 3 broad categories: nonspecific low back pain, back pain potentially associated with radiculopathy or spinal stenosis, or back pain potentially associated with another specific spinal cause. The history should include assessment of psychosocial risk factors, which predict risk for chronic disabling back pain (strong recommendation, moderate-quality evidence).

Recommendation 2: Clinicians should not routinely obtain imaging or other diagnostic tests in patients with nonspecific low back pain (strong recommendation, moderate-quality evidence).

Recommendation 3: Clinicians should perform diagnostic imaging and testing for patients with low back pain when severe or progressive neurologic deficits are present or when serious underlying conditions are suspected on the basis of history and physical examination (strong recommendation, moderate-quality evidence).

Recommendation 4: Clinicians should evaluate patients with persistent low back pain and signs or symptoms of radiculopathy or spinal stenosis with magnetic resonance imaging (preferred) or computed tomography only if they are potential candidates for surgery or epidural steroid injection (for suspected radiculopathy) (strong recommendation, moderate-quality evidence).

Recommendation 5: Clinicians should provide patients with evidence-based information on low back pain with regard to their expected course, advise patients to remain active, and provide information about effective self-care options (strong recommendation, moderate-quality evidence).


« Previous | Next Article »
Table of Contents

This Article

October 2, 2007
vol. 147 no. 7 478-491

Abstract [FREE](#)
Figures [FREE](#)
» Full Text [FREE](#)
Full Text (PDF) [FREE](#)
Audio Summary
Video News Release
Summary for Patients

Services

-  Related Clinical Content
- E-mail this article
- Alert me when new comments are published
- Similar articles in Annals
- Similar articles in PubMed
- Add to Personal Archive
- Download to citation manager
- Permissions for Course Packs

Comments

Submit a comment
Read comments (6)

- + Citing Articles
- + Google Scholar
- + PubMed
- + Related Content
- + Social Bookmarking

Navigate This Article

Top
Abstract
Methods
Recommendations: Evaluation of Low Back Pain
Recommendations: Treatment of Low Back Pain
Glossary
Article and Author Information
References

ACP AMERICAN COLLEGE OF PHYSICIANS®
INTERNAL MEDICINE | Doctors for Adults

Current Issue

March 15, 2011, 154 (6)

Alert me to current issues

in the clinic
ACP Journal Club®
ACP Journal Club®
PLUS

INFORMATION for:

Authors
Reviewers
Press
Readers
Subscribers
Institutions/Libraries/Agencies
Advertisers

► Most Read
► Most Commented On
► Annals in the News
► Clinical Trials
► Comparative Effectiveness

SERVICES and TOOLS:

iPhone/iPad App
Career Connection
Patient Information
CME
Teaching Tools
Feedback



Recommendation 6: For patients with low back pain, clinicians should consider the use of medications with proven benefits in conjunction with back care information and self-care. Clinicians should assess severity of baseline pain and functional deficits, potential benefits, risks, and relative lack of long-term efficacy and safety data before initiating therapy (strong recommendation, moderate-quality evidence). For most patients, first-line medication options are acetaminophen or nonsteroidal anti-inflammatory drugs.

Recommendation 7: For patients who do not improve with self-care options, clinicians should consider the addition of nonpharmacologic therapy with proven benefits—for acute low back pain, spinal manipulation; for chronic or subacute low back pain, intensive interdisciplinary rehabilitation, exercise therapy, acupuncture, massage therapy, spinal manipulation, yoga, cognitive-behavioral therapy, or progressive relaxation (weak recommendation, moderate-quality evidence).

* This paper, written by Roger Chou, MD; Amir Qaseem, MD, PhD, MHA; Vincenza Snow, MD; Donald Casey, MD, MPH, MBA; J. Thomas Cross Jr., MD, MPH; Paul Shekelle, MD, PhD; and Douglas K. Owens, MD, MS, was developed for the American College of Physicians' Clinical Efficacy Assessment Subcommittee and the American College of Physicians/American Pain Society Low Back Pain Guidelines Panel. For members of these groups, see end of text. Approved by the American College of Physicians Board of Regents on 14 July 2007. Approved by the American Pain Society Board Executive Committee on 18 July 2007.

Low back pain is the fifth most common reason for all physician visits in the United States (1, 2). Approximately one quarter of U.S. adults reported having low back pain lasting at least 1 whole day in the past 3 months (2), and 7.6% reported at least 1 episode of severe acute low back pain (see Glossary) within a 1-year period (3). Low back pain is also very costly: Total incremental direct health care costs attributable to low back pain in the U.S. were estimated at \$26.3 billion in 1998 (4). In addition, indirect costs related to days lost from work are substantial, with approximately 2% of the U.S. work force compensated for back injuries each year (5).

Many patients have self-limited episodes of acute low back pain and do not seek medical care (3). Among those who do seek medical care, pain, disability, and return to work typically improve rapidly in the first month (6). However, up to one third of patients report persistent back pain of at least moderate intensity 1 year after an acute episode, and 1 in 5 report substantial limitations in activity (7). Approximately 5% of the people with back pain disability account for 75% of the costs associated with low back pain (8).

Many options are available for evaluation and management of low back pain. However, there has been little consensus, either within or between specialties, on appropriate clinical evaluation (9) and management (10) of low back pain. Numerous studies show unexplained, large variations in use of diagnostic tests and treatments (11, 12). Despite wide variations in practice, patients seem to experience broadly similar outcomes, although costs of care can differ substantially among and within specialties (13, 14).

The purpose of this guideline is to present the available evidence for evaluation and management of acute and chronic low back pain (see Glossary) in primary care settings. The target audience for this guideline is all clinicians caring for patients with low (lumbar) back pain of any duration, either with or without leg pain. The target patient population is adults with acute and chronic low back pain not associated with major trauma. Children or adolescents with low back pain; pregnant women; and patients with low back pain from sources outside the back (nonspinal low back pain), fibromyalgia or other myofascial pain syndromes, and thoracic or cervical back pain are not included. These recommendations are based on a systematic evidence review summarized in 2 background papers by Chou and colleagues in this issue (15, 16) from an evidence report by the American Pain Society (17). The evidence report (17) discusses the evidence for the evaluation, and the 2 background papers (15, 16) summarize the evidence for management.



> FIND OUT MORE

Advertisement

Give yourself credit...

My CME
from *Annals of Internal Medicine*



- Take 20 CME quizzes a year for FREE
- View your completed courses, print your certificates, and more with My CME
- New CME quizzes are added with each new issue of *Annals*

Straightforward. Uncomplicated.

Free CME is an individual subscriber benefit.

VIEWED 03-16-2011

Methods

The literature search for this guideline included studies from MEDLINE (1966 through November 2006), the Cochrane Database of Systematic Reviews, the Cochrane Central Register of Controlled Trials, and EMBASE. The literature search included all English-language articles reporting on randomized, controlled trials of nonpregnant adults (age >18 years) with low back pain (alone or with leg pain) of any duration that evaluated a target medication and reported at least 1 of the following outcomes: back-specific function, generic health status, pain, work disability, or patient satisfaction. The American College of Physicians (ACP) and the American Pain Society (APS) convened a multidisciplinary panel of experts to develop the key questions and scope used to guide the evidence report, review its results, and formulate recommendations. The background papers by Chou and colleagues (15, 16) provide details about the methods used for the systematic evidence review.

This guideline grades its recommendations by using the ACP's clinical practice guidelines grading system, adapted from the classification developed by the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) work group (Appendix Table 1) (18). The evidence in this guideline was first evaluated by the ACP/APS panel by using a system adopted from the U.S. Preventive Services Task Force for grading strength of evidence, estimating magnitude of benefits, and assigning summary ratings (Appendix Tables 2, 3, and 4) (19). The evidence was independently reviewed by the ACP's Clinical Efficacy Assessment Subcommittee. The ratings for individual low back pain interventions discussed in this guideline are summarized in Appendix Table 5 for acute low back pain (<4 weeks' duration) and in Appendix Table 6 for chronic/subacute low back pain (>4 weeks' duration). This guideline considered interventions to have “proven” benefits only when they were supported by at least fair-quality evidence and were associated with at least moderate benefits (or small benefits but no significant harms, costs, or burdens). Figures 1 and 2 present an accompanying algorithm.

VIEWED 03-16-2011

View this table:

» In this window » In a new window

Appendix Table 1. The American College of Physicians Clinical Practice Guidelines Grading System

View this table:

» In this window » In a new window

Appendix Table 2. Methods for Grading the Strength of the Overall Evidence for an Intervention

View this table:

» In this window » In a new window

Appendix Table 3. Definitions for Estimating Magnitude of Effects

View this table:

» In this window » In a new window

Appendix Table 4. Recommendations and Summary Ratings

View this table:

» In this window » In a new window

Appendix Table 5. Level of Evidence and Summary Grades for Noninvasive Interventions in Patients with Acute Low Back Pain

View this table:

» In this window » In a new window


Appendix Table 6. Level of Evidence and Summary Grades for Noninvasive Interventions in Patients with Chronic or Subacute Low Back Pain



View larger version:

» In this page » In a new window
» Download as PowerPoint Slide

Figure 1.). Do not use this algorithm for back pain associated with major trauma, nonspinal back pain, or back pain due to systemic illness. CRP = C-reactive protein; EMG = electromyography; ESR = erythrocyte sedimentation rate; MRI = magnetic resonance imaging; NCV = nerve conduction velocity. Initial evaluation of low back pain (LBP)



View larger version:

» In this page » In a new window
» Download as PowerPoint Slide

Figure 2.). MRI = magnetic resonance imaging; NSAIDs = nonsteroidal anti-inflammatory drugs; TCA = tricyclic antidepressants. Management of low back pain (LBP)

Recommendations: Evaluation of Low Back Pain

Recommendation 1: Clinicians should conduct a focused history and physical examination to help place patients with low back pain into 1 of 3 broad categories: nonspecific low back pain, back pain potentially associated with radiculopathy or spinal stenosis, or back pain potentially associated with another specific spinal cause. The history should include assessment of psychosocial risk factors, which predict risk for chronic disabling back pain (strong recommendation, moderate-quality evidence).

More than 85% of patients who present to primary care have low back pain that cannot reliably be attributed to a specific disease or spinal abnormality (nonspecific low back pain [see Glossary]) (20). Attempts to identify specific anatomical sources of low back pain in such patients have not been validated in rigorous studies, and classification schemes frequently conflict with one another (21). Moreover, no evidence suggests that labeling most patients with low back pain by using specific anatomical diagnoses improves outcomes. In a minority of

VIEWED 03-16-2011

patients presenting for initial evaluation in a primary care setting, low back pain is caused by a specific disorder, such as cancer (approximately 0.7% of cases), compression fracture (4%), or spinal infection (0.01%) (22). Estimates for prevalence of ankylosing spondylitis in primary care patients range from 0.3% (22) to 5% (23). Spinal stenosis (see [Glossary](#)) and symptomatic herniated disc (see [Glossary](#)) are present in about 3% and 4% of patients, respectively. The cauda equina syndrome (see [Glossary](#)) is most commonly associated with massive midline disc herniation but is rare, with an estimated prevalence of 0.04% among patients with low back pain (24).

A practical approach to assessment is to do a focused history and physical examination to determine the likelihood of specific underlying conditions and measure the presence and level of neurologic involvement (24, 25). Such an approach facilitates classification of patients into 1 of 3 broad categories: nonspecific low back pain, back pain potentially associated with radiculopathy (see [Glossary](#)) or spinal stenosis (suggested by the presence of sciatica [see [Glossary](#)] or pseudoclaudication), and back pain potentially associated with another specific spinal cause. The latter category includes the small proportion of patients with serious or progressive neurologic deficits or underlying conditions requiring prompt evaluation (such as tumor, infection, or the cauda equina syndrome), as well as patients with other conditions that may respond to specific treatments (such as ankylosing spondylitis or vertebral compression fracture).

Diagnostic triage into 1 of these 3 categories helps guide subsequent decision making. Clinicians should inquire about the location of pain, frequency of symptoms, and duration of pain, as well as any history of previous symptoms, treatment, and response to treatment. The possibility of low back pain due to problems outside the back, such as pancreatitis, nephrolithiasis, or aortic aneurysm, or systemic illnesses, such as endocarditis or viral syndromes, should be considered. All patients should be evaluated for the presence of rapidly progressive or severe neurologic deficits, including motor deficits at more than 1 level, fecal incontinence, and bladder dysfunction. The most frequent finding in the cauda equina syndrome is urinary retention (97% sensitivity) (24). In patients without urinary retention, the probability of the cauda equina syndrome is approximately 1 in 10 000.

Clinicians should also ask about risk factors for cancer and infection. In a large, prospective study from a primary care setting, a history of cancer (positive likelihood ratio, 14.7), unexplained weight loss (positive likelihood ratio, 2.7), failure to improve after 1 month (positive likelihood ratio, 3.0), and age older than 50 years (positive likelihood ratio, 2.7) were each associated with a higher likelihood for cancer (26). The posttest probability of cancer in patients presenting with back pain increases from approximately 0.7% to 9% in patients with a history of cancer (not including nonmelanoma skin cancer). In patients with any 1 of the other 3 risk factors, the likelihood of cancer only increases to approximately 1.2% (26). Features predicting the presence of vertebral infection have not been well studied but may include fever, intravenous drug use, or recent infection (22). Clinicians should also consider risk factors for vertebral compression fracture, such as older age, history of osteoporosis, and steroid use, and ankylosing spondylitis, such as younger age, morning stiffness, improvement with exercise (see [Glossary](#)), alternating buttock pain, and awakening due to back pain during the second part of the night only (27), as specific treatments are available for these conditions. Clinicians should be aware that criteria for diagnosing early ankylosing spondylitis (before the development of radiographic abnormalities) are evolving (28).

In patients with back and leg pain, a typical history for sciatica (back and leg pain in a typical lumbar nerve root distribution) has a fairly high sensitivity but uncertain specificity for herniated disc (29, 30). More than 90% of symptomatic lumbar disc herniations (back and leg pain due to a prolapsed lumbar disc compressing a nerve root) occur at the L4/L5 and L5/S1 levels. A focused examination that includes straight-leg-raise testing (see [Glossary](#)) and a neurologic examination that includes evaluation of knee strength and reflexes (L4 nerve root), great toe and foot dorsiflexion strength (L5 nerve root), foot

VIEWED 03-16-2011

plantarflexion and ankle reflexes (S1 nerve root), and distribution of sensory symptoms should be done to assess the presence and severity of nerve root dysfunction. A positive result on the straight-leg-raise test (defined as reproduction of the patient's sciatica between 30 and 70 degrees of leg elevation) (24) has a relatively high sensitivity (91% [95% CI, 82% to 94%]) but modest specificity (26% [CI, 16% to 38%]) for diagnosing herniated disc (31). By contrast, the crossed straight-leg-raise test is more specific (88% [CI, 86% to 90%]) but less sensitive (29% [CI, 24% to 34%]).

Evidence on the utility of history and examination for identifying lumbar spinal stenosis is sparse (32). High-quality studies showed a trade-off between sensitivities and specificities, resulting in modest or poor positive likelihood ratios (1.2 for pseudoclaudication and 2.2 for radiating leg pain) (32). Changing symptoms on downhill treadmill testing are associated with the highest positive likelihood ratio (3.1). The usefulness of pain relieved by sitting for predicting presence of spinal stenosis ranges from poor to high (32). Age older than 65 years was associated with a positive likelihood ratio of 2.5 and a negative likelihood ratio of 0.33 in 1 lower-quality study (33). Other findings have only been evaluated in lower-quality studies or are poorly predictive for lumbar spinal stenosis.

Psychosocial factors and emotional distress should be assessed because they are stronger predictors of low back pain outcomes than either physical examination findings or severity and duration of pain (6, 34, 35). Assessment of psychosocial factors identifies patients who may have delayed recovery and could help target interventions, as 1 trial in a referral setting found intensive multidisciplinary rehabilitation more effective than usual care in patients with acute or subacute low back pain identified as having risk factors for chronic back pain disability (36). Direct evidence on effective primary care interventions for identifying and treating such factors in patients with acute low back pain is lacking (37, 38), although this is an area of active research. Evidence is currently insufficient to recommend optimal methods for assessing psychosocial factors and emotional distress. However, psychosocial factors that may predict poorer low back pain outcomes include presence of depression, passive coping strategies, job dissatisfaction, higher disability levels, disputed compensation claims, or somatization (34, 35, 39).

Evidence is also insufficient to guide appropriate intervals or methods (such as office visit vs. telephone follow-up) for reassessment of history, physical examination, or psychosocial factors. However, patients with acute low back pain generally experience substantial improvement in the first month after initial presentation (6, 40), suggesting that a reasonable approach is to reevaluate patients with persistent, unimproved symptoms after 1 month. In patients with severe pain or functional deficits, older patients, or patients with signs of radiculopathy or spinal stenosis (see recommendation 4), earlier or more frequent reevaluation may also be appropriate.

Recommendation 2: Clinicians should not routinely obtain imaging or other diagnostic tests in patients with nonspecific low back pain (strong recommendation, moderate-quality evidence).

There is no evidence that routine plain radiography in patients with nonspecific low back pain is associated with a greater improvement in patient outcomes than selective imaging (41–43). In addition, exposure to unnecessary ionizing radiation should be avoided. This issue is of particular concern in young women because the amount of gonadal radiation from obtaining a single plain radiograph (2 views) of the lumbar spine is equivalent to being exposed to a daily chest radiograph for more than 1 year (44). Routine advanced imaging (computed tomography [CT] or magnetic resonance imaging [MRI]) is also not associated with improved patient outcomes (45) and identifies many radiographic abnormalities that are poorly correlated with symptoms (22) but could lead to additional, possibly unnecessary interventions (46, 47).

Plain radiography is recommended for initial evaluation of possible vertebral compression fracture in selected higher-risk patients, such as those with a history of osteoporosis or steroid use (22). Evidence to guide optimal imaging

VIEWED 03-16-2011

strategies is not available for low back pain that persists for more than 1 to 2 months despite standard therapies if there are no symptoms suggesting radiculopathy or spinal stenosis, although plain radiography may be a reasonable initial option (see recommendation 4 for imaging recommendations in patients with symptoms suggesting radiculopathy or spinal stenosis). Thermography and electrophysiologic testing are not recommended for evaluation of nonspecific low back pain.

Recommendation 3: Clinicians should perform diagnostic imaging and testing for patients with low back pain when severe or progressive neurologic deficits are present or when serious underlying conditions are suspected on the basis of history and physical examination (strong recommendation, moderate-quality evidence).

Prompt work-up with MRI or CT is recommended in patients who have severe or progressive neurologic deficits or are suspected of having a serious underlying condition (such as vertebral infection, the cauda equina syndrome, or cancer with impending spinal cord compression) because delayed diagnosis and treatment are associated with poorer outcomes (48–50). Magnetic resonance imaging is generally preferred over CT if available because it does not use ionizing radiation and provides better visualization of soft tissue, vertebral marrow, and the spinal canal (22). There is insufficient evidence to guide precise recommendations on diagnostic strategies in patients who have risk factors for cancer but no signs of spinal cord compression. Several strategies have been proposed for such patients (22, 51), but none have been prospectively evaluated. Proposed strategies generally recommend plain radiography or measurement of erythrocyte sedimentation rate (a rate ≥ 20 mm/h is associated with 78% sensitivity and 67% specificity for cancer [29]), with MRI reserved for patients with abnormalities on initial testing (22, 51). An alternative strategy is to directly perform MRI in patients with a history of cancer, the strongest predictor of vertebral cancer (51). For patients older than 50 years of age without other risk factors for cancer, delaying imaging while offering standard treatments and reevaluating within 1 month may also be a reasonable option (52).

Recommendation 4: Clinicians should evaluate patients with persistent low back pain and signs or symptoms of radiculopathy or spinal stenosis with MRI (preferred) or CT only if they are potential candidates for surgery or epidural steroid injection (for suspected radiculopathy) (strong recommendation, moderate-quality evidence).

The natural history of lumbar disc herniation with radiculopathy in most patients is for improvement within the first 4 weeks with noninvasive management (53, 54). There is no compelling evidence that routine imaging affects treatment decisions or improves outcomes (55). For prolapsed lumbar disc with persistent radicular symptoms despite noninvasive therapy, discectomy or epidural steroids are potential treatment options (56–60). Surgery is also a treatment option for persistent symptoms associated with spinal stenosis (61–64).

Magnetic resonance imaging (preferred if available) or CT is recommended for evaluating patients with persistent back and leg pain who are potential candidates for invasive interventions—plain radiography cannot visualize discs or accurately evaluate the degree of spinal stenosis (22). However, clinicians should be aware that findings on MRI or CT (such as bulging disc without nerve root impingement) are often nonspecific. Recommendations for specific invasive interventions, interpretation of radiographic findings, and additional work-up (such as electrophysiologic testing) are beyond the scope of this guideline, but decisions should be based on the clinical correlation between symptoms and radiographic findings, severity of symptoms, patient preferences, surgical risks (including the patient's comorbid conditions), and costs and will generally require specialist input.

Recommendations: Treatment of Low Back Pain

Recommendation 5: Clinicians should provide patients with evidence-based information on low back pain with regard to their expected course, advise

VIEWED 03-16-2011

patients to remain active, and provide information about effective self-care options (strong recommendation, moderate-quality evidence).

Clinicians should inform all patients of the generally favorable prognosis of acute low back pain with or without sciatica, including a high likelihood for substantial improvement in the first month (6, 40). Clinicians should explain that early, routine imaging and other tests usually cannot identify a precise cause, do not improve patient outcomes, and incur additional expenses. Clinicians should also review indications for reassessment and diagnostic testing (see recommendations 1 and 4). General advice on self-management for nonspecific low back pain should include recommendations to remain active, which is more effective than resting in bed for patients with acute or subacute low back pain (65, 66). If patients require periods of bed rest to relieve severe symptoms, they should be encouraged to return to normal activities as soon as possible. Self-care education books (see Glossary) based on evidence-based guidelines, such as *The Back Book*(67), are recommended because they are an inexpensive and efficient method for supplementing clinician-provided back information and advice and are similar or only slightly inferior in effectiveness to such costlier interventions as supervised exercise therapy, acupuncture (see Glossary), massage (see Glossary), and spinal manipulation (see Glossary) (65, 66, 68–70). Other methods for providing self-care education, such as e-mail discussion groups, layperson-led groups, videos, and group classes, are not as well studied.

Factors to consider when giving advice about activity limitations to workers with low back pain are the patient's age and general health and the physical demands of required job tasks. However, evidence is insufficient to guide specific recommendations about the utility of modified work for facilitating return to work (71). For worker's compensation claims, clinicians should refer to specific regulations for their area of practice, as rules vary substantially from state to state. Brief individualized educational interventions (defined as a detailed clinical examination and advice, typically lasting several hours over 1 to 2 sessions) (see Glossary) can reduce sick leave in workers with subacute low back pain (72–74).

Application of heat by heating pads or heated blankets is a self-care option (see Glossary) for short-term relief of acute low back pain (75). In patients with chronic low back pain, firm mattresses are less likely than a medium-firm mattress to lead to improvement (76). There is insufficient evidence to recommend lumbar supports (77) or the application of cold packs (75) as self-care options.

Although evidence is insufficient to guide specific self-management recommendations for patients with acute radiculopathy or spinal stenosis, some trials enrolled mixed populations of patients with and without sciatica, suggesting that applying principles similar to those used for nonspecific low back pain is a reasonable approach (see also recommendation 4).

Recommendation 6: For patients with low back pain, clinicians should consider the use of medications with proven benefits in conjunction with back care information and self-care. Clinicians should assess severity of baseline pain and functional deficits, potential benefits, risks, and relative lack of long-term efficacy and safety data before initiating therapy (strong recommendation, moderate-quality evidence). For most patients, first-line medication options are acetaminophen or nonsteroidal anti-inflammatory drugs (NSAIDs).

Medications in several classes have been shown to have moderate, primarily short-term benefits for patients with low back pain. Each class of medication is associated with unique trade-offs involving benefits, risks, and costs. For example, acetaminophen is a slightly weaker analgesic than NSAIDs (<10 points on a 100-point visual analogue pain scale) (78–82) but is a reasonable first-line option for treatment of acute or chronic low back pain because of a more favorable safety profile and low cost (79, 82–84). However, acetaminophen is associated with asymptomatic elevations of aminotransferase levels at dosages of 4 g/d (the upper limit of U.S. Food and Drug Administration-[FDA] approved dosing) even in healthy adults, although the clinical significance of these findings are uncertain (85). Nonselective NSAIDs are more effective for pain

VIEWED 03-16-2011

relief than is acetaminophen (80), but they are associated with well-known gastrointestinal and renovascular risks (83). In addition, there is an association between exposure to cyclooxygenase-2-selective or most nonselective NSAIDs and increased risk for myocardial infarction (86). Clinicians should therefore assess cardiovascular and gastrointestinal risk factors before prescribing NSAIDs and recommend the lowest effective doses for the shortest periods necessary. Clinicians should also remain alert for new evidence about which NSAIDs are safest and consider strategies for minimizing adverse events in higher-risk patients who are prescribed NSAIDs (such as co-administration with a proton-pump inhibitor) (87). There is insufficient evidence to recommend for or against analgesic doses of aspirin in patients with low back pain (88).

Opioid analgesics or tramadol are an option when used judiciously in patients with acute or chronic low back pain who have severe, disabling pain that is not controlled (or is unlikely to be controlled) with acetaminophen and NSAIDs. Because of substantial risks, including aberrant drug-related behaviors with long-term use in patients vulnerable or potentially vulnerable to abuse or addiction, potential benefits and harms of opioid analgesics should be carefully weighed before starting therapy (89–91). Failure to respond to a time-limited course of opioids should lead to reassessment and consideration of alternative therapies or referral for further evaluation (92–94). Evidence is insufficient to recommend one opioid over another (95).

The [Glossary](#) term *skeletal muscle relaxants* refers to a diverse group of medications, some with unclear mechanisms of action, grouped together because they carry FDA-approved indications for treatment of musculoskeletal conditions or spasticity. Although the antispasticity drug tizanidine has been well studied for low back pain, there is little evidence for the efficacy of baclofen or dantrolene, the other FDA-approved drugs for the treatment of spasticity (96). Other medications in the skeletal muscle relaxant class are an option for short-term relief of acute low back pain, but all are associated with central nervous system adverse effects (primarily sedation). There is no compelling evidence that skeletal muscle relaxants differ in efficacy or safety (96, 97). Because skeletal muscle relaxants are not pharmacologically related, however, risk-benefit profiles could in theory vary substantially. For example, carisoprodol is metabolized to meprobamate (a medication associated with risks for abuse and overdose), dantrolene carries a black box warning for potentially fatal hepatotoxicity, and both tizanidine and chlorzoxazone are associated with hepatotoxicity that is generally reversible and usually not serious.

Tricyclic antidepressants are an option for pain relief in patients with chronic low back pain and no contraindications to this class of medications (98, 99). Antidepressants in the selective serotonin reuptake inhibitor class and trazodone have not been shown to be effective for low back pain, and serotonin-norepinephrine reuptake inhibitors (duloxetine and venlafaxine) have not yet been evaluated for low back pain. Clinicians should bear in mind, however, that depression is common in patients with chronic low back pain and should be assessed and treated appropriately (100).

Gabapentin is associated with small, short-term benefits in patients with radiculopathy (101, 102) and has not been directly compared with other medications or treatments. There is insufficient evidence to recommend for or against other antiepileptic drugs for back pain with or without radiculopathy. For acute or chronic low back pain, benzodiazepines seem similarly effective to skeletal muscle relaxants for short-term pain relief (96) but are also associated with risks for abuse, addiction, and tolerance. Neither benzodiazepines nor gabapentin are FDA-approved for treatment of low back pain (with or without radiculopathy). If a benzodiazepine is used, a time-limited course of therapy is recommended.

Herbal therapies, such as devil's claw, willow bark, and capsicum, seem to be safe options for acute exacerbations of chronic low back pain, but benefits range from small to moderate. In addition, many of the published trials were led by the same investigator, which could limit applicability of findings to other settings (103).

VIEWED 03-16-2011

Systemic corticosteroids are not recommended for treatment of low back pain with or without sciatica, because they have not been shown to be more effective than placebo (104–107).

Most medication trials evaluated patients with nonspecific low back pain or mixed populations with and without sciatica. There is little evidence to guide specific recommendations for medications (other than gabapentin) for patients with sciatica or spinal stenosis. Evidence is also limited on the benefits and risks associated with long-term use of medications for low back pain. Therefore, extended courses of medications should generally be reserved for patients clearly showing continued benefits from therapy without major adverse events.

Recommendation 7: For patients who do not improve with self-care options, clinicians should consider the addition of nonpharmacologic therapy with proven benefits—for acute low back pain, spinal manipulation; for chronic or subacute low back pain, intensive interdisciplinary rehabilitation, exercise therapy, acupuncture, massage therapy, spinal manipulation, yoga, cognitive-behavioral therapy, or progressive relaxation (weak recommendation, moderate-quality evidence).

For acute low back pain (duration <4 weeks), spinal manipulation administered by providers with appropriate training is associated with small to moderate short-term benefits (108). Supervised exercise therapy and home exercise regimens are not effective for acute low back pain (109), and the optimal time to start exercise therapy after the onset of symptoms is unclear. Other guidelines suggest starting exercise after 2 to 6 weeks, but these recommendations seem to be based on poor-quality evidence (25, 110). Other nonpharmacologic treatments have not been proven to be effective for acute low back pain.

For subacute (duration >4 to 8 weeks) low back pain, intensive interdisciplinary rehabilitation (defined as an intervention that includes a physician consultation coordinated with a psychological, physical therapy, social, or vocational intervention) (see Glossary) is moderately effective (111), and functional restoration (see Glossary) with a cognitive-behavioral component reduces work absenteeism due to low back pain in occupational settings (112). There is little evidence on effectiveness of other treatments specifically for subacute low back pain (113). However, many trials enrolled mixed populations of patients with chronic and subacute symptoms, suggesting that results may reasonably be applied to both situations.

For chronic low back pain, moderately effective nonpharmacologic therapies include acupuncture (114, 115), exercise therapy (109), massage therapy (116), Viniyoga-style yoga (see Glossary) (70), cognitive-behavioral therapy or progressive relaxation (see Glossary) (117, 118), spinal manipulation (108), and intensive interdisciplinary rehabilitation (119), although the level of supporting evidence for different therapies varies from fair to good (Appendix Table 6). In meta-regression analyses, exercise programs that incorporate individual tailoring, supervision, stretching, and strengthening are associated with the best outcomes (109). The evidence is insufficient to conclude that benefits of manipulation vary according to the profession of the manipulator (chiropractor vs. other clinician trained in manipulation) or according to presence or absence of radiating pain (108). With the exception of continuous or intermittent traction (see Glossary), which has not been shown to be effective in patients with sciatica (120–122), few trials have evaluated the effectiveness of treatments specifically in patients with radicular pain (122) or symptoms of spinal stenosis. In addition, there is insufficient evidence to recommend any specific treatment as first-line therapy. Patient expectations of benefit from a treatment should be considered in choosing interventions because they seem to influence outcomes (123). Some interventions (such as intensive interdisciplinary rehabilitation) may not be available in all settings, and costs for similarly effective interventions can vary substantially. There is insufficient evidence to recommend the use of decision tools or other methods for tailoring therapy in primary care, although initial data are promising (124–126).

Transcutaneous electrical nerve stimulation (see Glossary) and intermittent or continuous traction (in patients with or without sciatica) have not been proven

VIEWED 03-16-2011

effective for chronic low back pain (Appendix Table 6). Acupressure (see Glossary), neuroreflexotherapy (see Glossary), and spa therapy (see Glossary) have not been studied in the United States, and percutaneous electrical nerve stimulation (see Glossary) is not widely available. There is insufficient evidence to recommend interferential therapy (see Glossary), low-level laser therapy (see Glossary), shortwave diathermy (see Glossary), or ultrasonography. Evidence is inconsistent on back schools (see Glossary), which have primarily been evaluated in occupational settings, with some trials showing small, short-term benefits (127).

It may be appropriate to consider consultation with a back specialist when patients with nonspecific low back pain do not respond to standard noninvasive therapies. However, there is insufficient evidence to guide specific recommendations on the timing of or indications for referral, and expertise in management of low back pain varies substantially among clinicians from different disciplines (including primary care providers). In general, decisions about consultation should be individualized and based on assessments of patient symptoms and response to interventions, the experience and training of the primary care clinician, and the availability of specialists with relevant expertise. In considering referral for possible surgery or other invasive interventions, other published guidelines suggest referring patients with nonspecific low back pain after a minimum of 3 months (25) to 2 years (128) of failed nonsurgical interventions. Although specific suggestions about timing of referral are somewhat arbitrary, one factor to consider is that trials of surgery for nonspecific low back pain included only patients with at least 1 year of symptoms (129–131). Other recommendations for invasive interventions are addressed in a separate guideline from the APS (17).

Glossary

General

Acute low back pain: Low back pain present for fewer than 4 weeks, sometimes grouped with subacute low back pain as symptoms present for fewer than 3 months.

Cauda equina syndrome: Compression on nerve roots from the lower cord segments, usually due to a massive, centrally herniated disc, which can result in urinary retention or incontinence from loss of sphincter function, bilateral motor weakness of the lower extremities, and saddle anesthesia.

Chronic low back pain: Low back pain present for more than 3 months.

Herniated disc: Herniation of the nucleus pulposus of an intervertebral disc through its fibrous outer covering, which can result in compression of adjacent nerve roots or other structures.

Neurogenic claudication: Symptoms of leg pain (and occasionally weakness) on walking or standing, relieved by sitting or spinal flexion, associated with spinal stenosis.

Nonspecific low back pain: Pain occurring primarily in the back with no signs of a serious underlying condition (such as cancer, infection, or cauda equina syndrome), spinal stenosis or radiculopathy, or another specific spinal cause (such as vertebral compression fracture or ankylosing spondylitis). Degenerative changes on lumbar imaging are usually considered nonspecific, as they correlate poorly with symptoms.

Radiculopathy: Dysfunction of a nerve root associated with pain, sensory impairment, weakness, or diminished deep tendon reflexes in a nerve root distribution.

Sciatica: Pain radiating down the leg below the knee in the distribution of the sciatic nerve, suggesting nerve root compromise due to mechanical pressure or inflammation. Sciatica is the most common symptom of lumbar radiculopathy.

Spinal stenosis: Narrowing of the spinal canal that may result in bony

VIEWED 03-16-2011

constriction of the cauda equina and the emerging nerve roots.

Straight-leg-raise test: A procedure in which the hip is flexed with the knee extended in order to passively stretch the sciatic nerve and elicit symptoms suggesting nerve root tension. A positive test is usually considered reproduction of the patient's sciatica when the leg is raised between 30 and 70 degrees. Reproduction of the patient's sciatica when the unaffected leg is lifted is referred to as a positive "crossed" straight-leg-raise test.

Interventions

Acupressure: An intervention consisting of manipulation with the fingers instead of needles at specific acupuncture points.

Acupuncture: An intervention consisting of the insertion of needles at specific acupuncture points.

Back school: An intervention consisting of education and a skills program, including exercise therapy, in which all lessons are given to groups of patients and supervised by a paramedical therapist or medical specialist.

Brief individualized educational interventions: Individualized assessment and education about low back pain problems without supervised exercise therapy or other specific interventions. As we defined them, brief educational interventions differ from back schools because they do not involve group education or supervised exercise.

Exercise: A supervised exercise program or formal home exercise regimen, ranging from programs aimed at general physical fitness or aerobic exercise to programs aimed at muscle strengthening, flexibility, stretching, or different combinations of these elements.

Functional restoration (also called physical conditioning, work hardening, or work conditioning): An intervention that involves simulated or actual work tests in a supervised environment in order to enhance job performance skills and improve strength, endurance, flexibility, and cardiovascular fitness in injured workers.

Interdisciplinary rehabilitation (also called multidisciplinary therapy): An intervention that combines and coordinates physical, vocational, and behavioral components and is provided by multiple health care professionals with different clinical backgrounds. The intensity and content of interdisciplinary therapy varies widely.

Interferential therapy: The superficial application of a medium-frequency alternating current modulated to produce low frequencies up to 150 Hz. It is thought to increase blood flow to tissues and provide pain relief and is considered more comfortable for patients than transcutaneous electrical nerve stimulation.

Low-level laser therapy: The superficial application of lasers at wavelengths between 632 and 904 nm to the skin in order to apply electromagnetic energy to soft tissue. Optimal treatment parameters (wavelength, dosage, dose-intensity, and type of laser) are uncertain.

Massage: Soft tissue manipulation using the hands or a mechanical device through a variety of specific methods. The pressure and intensity used in different massage techniques vary widely.

Neuroreflexotherapy: A technique from Spain characterized by the temporary implantation of staples superficially into the skin over trigger points in the back and referred tender points in the ear. Neuroreflexotherapy is believed to stimulate different zones of the skin than acupuncture.

Percutaneous electrical nerve stimulation (PENS): An intervention that involves inserting acupuncture-like needles and applying low-level electrical stimulation. It differs from electroacupuncture in that the insertion points target dermatomal levels for local pathology, rather than acupuncture points. However, there is

VIEWED 03-16-2011

some uncertainty over whether PENS should be considered a novel therapy or a form of electroacupuncture.

Progressive relaxation: A technique which involves the deliberate tensing and relaxation of muscles, in order to facilitate the recognition and release of muscle tension.

Self-care options: Interventions that can be readily implemented by patients without seeing a clinician or that can be implemented on the basis of advice provided at a routine clinic visit.

Self-care education book: Reading material (books, booklets, or leaflets) that provide education and self-care advice for patients with low back pain. Although the specific content varies, self-care books are generally based on principles from published clinical practice guidelines and encourage a return to normal activity, adoption of a fitness program, and appropriate lifestyle modification, and they provide advice on coping strategies and managing flares.

Shortwave diathermy: Therapeutic elevation of the temperature of deep tissues by application of short-wave electromagnetic radiation with a frequency range from 10–100 MHz.

Spa therapy: An intervention involving several interventions, including mineral water bathing, usually with heated water, typically while staying at a spa resort.

Spinal manipulation: Manual therapy in which loads are applied to the spine by using short- or long-lever methods and high-velocity thrusts are applied to a spinal joint beyond its restricted range of movement. Spinal mobilization, or low-velocity, passive movements within or at the limit of joint range, is often used in conjunction with spinal manipulation.

Traction: An intervention involving drawing or pulling in order to stretch the lumbar spine. Various methods are used, usually involving a harness around the lower rib cage and the iliac crest, with the pulling action done by using free weights and a pulley, motorized equipment, inversion techniques, or an overhead harness.

Transcutaneous electrical nerve stimulation (TENS): Use of a small, battery-operated device to provide continuous electrical impulses via surface electrodes, with the goal of providing symptomatic relief by modifying pain perception.

Yoga: An intervention distinguished from traditional exercise therapy by the use of specific body positions, breathing techniques, and an emphasis on mental focus. Many styles of yoga are practiced, each emphasizing different postures and techniques.

Article and Author Information

Clinical Efficacy Assessment Subcommittee of the American College of Physicians: Douglas K. Owens, MD, MS (Chair)†; Donald E. Casey Jr., MD, MPH, MBA‡; J. Thomas Cross Jr., MD, MPH‡; Paul Dallas, MD; Nancy C. Dolan, MD; Mary Ann Forciea, MD; Lakshmi Halasyamani, MD; Robert H. Hopkins Jr., MD; and Paul Shekelle, MD, PhD‡. *Co-chairs and members of the American College of Physicians/American Pain Society Low Back Pain Guidelines Panel:* John D. Loeser, MD (Co-chair); Douglas K. Owens, MD, MS (Co-chair); Richard W. Rosenquist, MD (Co-chair); Paul M. Arnstein, RN, PhD, APRN-BC; Steven Julius Atlas, MD, MPH; Jamie Baisden, MD; Claire Bombardier, MD; Eugene J. Carragee, MD; John Anthony Carrino, MD, MPH; Donald E. Casey Jr., MD, MPH, MBA; Daniel Cherkin, PhD; Penney Cowan; J. Thomas Cross Jr., MD, MPH; Anthony Delitto, PhD, MHS; Robert J. Gatchel, Ph.D, ABPP; Lee Steven Glass, MD, JD; Martin Grabois, MD; Timothy R. Lubenow, MD; Kathryn Mueller, MD, MPH; Donald R. Murphy, DC, DACAN; Marco Pappagallo, MD; Kenneth G. Saag, MD, MSc; Paul G. Shekelle, MD, PhD; Steven P. Stanos, DO; and Eric Martin Wall, MD, MPH. *Participants from the Veterans Affairs/Department of Defense Evidence-Based Practice Workgroup:* Carla L. Cassidy, ANP, MSN; COL Leo L. Bennett, MC, MD, MPH;

VIEWED 03-16-2011

John Dooley, MD; LCDR Leslie Rassner, MD; Robert Ruff, MD, PhD; and Suzanne Ruff, MHCC. †Also a co-chair of the American College of Physicians/American Pain Society Low Back Pain Guidelines Panel. ‡Also members of the American College of Physicians/American Pain Society Low Back Pain Guidelines Panel.

Note: Clinical practice guidelines are “guides” only and may not apply to all patients and all clinical situations. Thus, they are not intended to override clinicians’ judgment. All ACP clinical practice guidelines are considered automatically withdrawn or invalid 5 years after publication or once an update has been issued.

Disclaimer: The authors of this article are responsible for its contents, including any clinical or treatment recommendations. The views and opinions expressed are those of Veterans Affairs/Department of Defense Evidence-Based Practice Workgroup members and do not necessarily reflect official Veterans Health Affairs or Department of Defense positions.

Acknowledgments: The authors thank Laurie Hoyt Huffman for reviewing the manuscript and providing helpful suggestions, Jayne Schablaske and Michelle Pappas for administrative assistance in preparing the manuscript, Andrew Hamilton for conducting the literature searches, and Oded Susskind for assistance in developing the algorithm.

Grant Support: Financial support for the development of this guideline comes exclusively from the ACP and APS operating budgets.

Potential Financial Conflicts of Interest: Honoraria: R. Chou (Bayer Healthcare Pharmaceuticals). Grants received: V. Snow (Centers for Disease Control and Prevention, Agency for Healthcare Research and Quality, Novo Nordisk, Pfizer Inc., Merck & Co. Inc., Bristol-Myers Squibb, Atlantic Philanthropies, Sanofi-Pasteur).

Requests for Single Reprints: Amir Qaseem, MD, PhD, MHA, American College of Physicians, 190 N. Independence Mall West, Philadelphia, PA 19106; e-mail, aqaseem@acponline.org.

Current Author Addresses: Dr. Chou: 3181 SW Sam Jackson Park Road, Mailcode BICC, Portland, OR 97239.

Drs. Qaseem and Snow: American College of Physicians, 190 N. Independence Mall West, Philadelphia, PA 19106.

Dr. Casey: 475 South Street, PO Box 1905, Morristown, NJ 07962.

Dr. Cross: 1761 South 8th Street, Suite H, Colorado Springs, CO 80906.

Dr. Shekelle: 1776 Main Street, Santa Monica, CA 90401.

Dr. Owens: 117 Encina Commons, Stanford, CA 94305

References

1. [] Hart LG, Deyo RA, Cherkin DC. Physician office visits for low back pain. Frequency, clinical evaluation, and treatment patterns from a U.S. national survey. Spine. 1995;20:11–9. [PMID: 7709270] » [Medline](#)
2. [] Deyo RA, Mirza SK, Martin BI. Back pain prevalence and visit rates: estimates from U.S. national surveys, 2002. Spine. 2006;31:2724–7. [PMID: 17077742] » [CrossRef](#) » [Medline](#)
3. [] Carey TS, Evans AT, Hadler NM, Lieberman G, Kalsbeek WD, Jackman AM, et al. Acute severe low back pain. A population-based study of prevalence and

» [CrossRef](#) » [Medline](#)

VIEWED 03-16-2011

care-seeking. Spine. 1996;21:339–44. [PMID: 8742211]

4. Luo X, Pietrobon R, Sun SX, Liu GG, Hey L. Estimates and patterns of direct health care expenditures among individuals with back pain in the United States. Spine. 2004;29:79–86. [PMID: 14699281] »CrossRef »Medline

5. Andersson GB. Epidemiological features of chronic low-back pain. Lancet. 1999;354:581–5. [PMID: 10470716] »CrossRef »Medline

6. Pengel LH, Herbert RD, Maher CG, Refshauge KM. Acute low back pain: systematic review of its prognosis. BMJ. 2003;327:323. [PMID: 12907487] »Abstract/FREE Full Text

7. Von Korff M, Saunders K. The course of back pain in primary care. Spine. 1996;21:2833–7; discussion 2838–9. [PMID: 9112707]

8. Frymoyer JW, Cats-Baril WL. An overview of the incidences and costs of low back pain. Orthop Clin North Am. 1991;22:263–71. [PMID: 1826550] »Medline

9. Cherkin DC, Deyo RA, Wheeler K, Ciol MA. Physician variation in diagnostic testing for low back pain. Who you see is what you get. Arthritis Rheum. 1994;37:15–22. [PMID: 8129759] »Medline

10. Cherkin DC, Deyo RA, Wheeler K, Ciol MA. Physician views about treating low back pain. The results of a national survey. Spine. 1995;20:1–9; discussion 9–10. [PMID: 7709266]

11. Cherkin DC, Deyo RA, Loeser JD, Bush T, Waddell G. An international comparison of back surgery rates. Spine. 1994;19:1201–6. [PMID: 8073310] »Medline

12. Volinn E, Mayer J, Diehr P, Van Koeveing D, Connell FA, Loeser JD. Small area analysis of surgery for low-back pain. Spine. 1992;17:575–81. [PMID: 1535726] »Medline

13. Carey TS, Garrett J, Jackman A, McLaughlin C, Fryer J, Smucker DR. The outcomes and costs of care for acute low back pain among patients seen by primary care practitioners, chiropractors, and orthopedic surgeons. The North Carolina Back Pain Project. N Engl J Med. 1995;333:913–7. [PMID: 7666878] »CrossRef »Medline

14. Shekelle PG, Markovich M, Louie R. Comparing the costs between provider types of episodes of back pain care. Spine. 1995;20:221–6; discussion 227. [PMID: 7716629]

15. Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline. Ann Intern Med. 2007;147:492–504.

16. Chou R, Huffman LH. Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. Ann Intern Med. 2007;147:505–14.

17. Chou R, Huffman L. Evaluation and management of low back pain: evidence review. Glenview, IL: American Pain Soc; 2007. [In press]

18. Guyatt G, Gutterman D, Baumann MH, Addrizzo-Harris D, Hylek EM, Phillips B, et al. Grading strength of recommendations and quality of evidence in clinical guidelines: report from an american college of chest physicians task force. Chest. 2006;129:174–81. [PMID: 16424429] »Abstract/FREE Full Text

19. Harris R, Helfand M, Woolf S, et.al. Methods Work Group, Third US Preventive Services Task Force. Current methods of the US Preventive Services Task Force: a review of the process. Am J Prev Med. 2001;20:21–35. [PMID: 11306229] »Medline

20. van Tulder MW, Assendelft WJ, Koes BW, Bouter LM. Spinal radiographic findings and nonspecific low back pain. A systematic review of observational studies. Spine. 1997;22:427–34. [PMID: 9055372] »CrossRef »Medline

21. Deyo RA. Practice variations, treatment fads, rising disability. Do we need a new clinical research paradigm? Spine. 1993;18:2153–62. [PMID: 8278825] »Medline

22. Jarvik JG, Deyo RA. Diagnostic evaluation of low back pain with emphasis on imaging. Ann Intern Med. 2002;137:586–97. [PMID: 12353946] »Abstract/FREE Full Text


23. Underwood MR, Dawes P. Inflammatory back pain in primary care. Br J Rheumatol. 1995;34:1074–7. [PMID: 8542211] »Abstract/FREE Full Text


24. Deyo RA, Rainville J, Kent DL. What can the history and physical examination tell us about low back pain? JAMA. 1992;268:760–5. [PMID: 1386391] »Abstract/FREE Full Text


25. Bigos S, Bowyer O, Braen G, Brown K, Deyo R, Haldeman S, et al. Acute Low Back Problems in Adults. Clinical Practice Guideline No. 14. AHCPR Publication No. 95–0642. Rockville, MD: Agency for Health Care Policy and


VIEWED 03-16-2011


Research, Public Health Service, U.S. Department of Health and Human Services; 1994.


26.  Deyo RA, Diehl AK. Cancer as a cause of back pain: frequency, clinical presentation, and diagnostic strategies. *J Gen Intern Med*. 1988;3:230–8. [PMID: 2967893] [» Medline](#)


27.  Rudwaleit M, Metter A, Listing J, Sieper J, Braun J. Inflammatory back pain in ankylosing spondylitis: a reassessment of the clinical history for application as classification and diagnostic criteria. *Arthritis Rheum*. 2006;54:569–78. [PMID: 16447233] [» CrossRef](#) [» Medline](#)


28.  Rudwaleit M, Khan MA, Sieper J. The challenge of diagnosis and classification in early ankylosing spondylitis: do we need new criteria? *Arthritis Rheum*. 2005;52:1000–8. [PMID: 15818678] [» CrossRef](#) [» Medline](#)


29.  van den Hoogen HM, Koes BW, van Eijk JT, Bouter LM. On the accuracy of history, physical examination, and erythrocyte sedimentation rate in diagnosing low back pain in general practice. A criteria-based review of the literature. *Spine*. 1995;20:318–27. [PMID: 7732468] [» Medline](#)


30.  Vroomen PC, de Krom MC, Knottnerus JA. Diagnostic value of history and physical examination in patients suspected of sciatica due to disc herniation: a systematic review. *J Neurol*. 1999;246:899–906. [PMID: 10552236] [» CrossRef](#) [» Medline](#)


31.  Devillé WL, van der Windt DA, Dzaferagić A, Bezemer PD, Bouter LM. The test of Lasègue: systematic review of the accuracy in diagnosing herniated discs. *Spine*. 2000;25:1140–7. [PMID: 10788860] [» CrossRef](#) [» Medline](#)


32.  de Graaf I, Prak A, Bierma-Zeinstra S, Thomas S, Peul W, Koes B. Diagnosis of lumbar spinal stenosis: a systematic review of the accuracy of diagnostic tests. *Spine*. 2006;31:1168–76. [PMID: 16648755] [» CrossRef](#) [» Medline](#)


33.  Katz JN, Dalgas M, Stucki G, Katz NP, Bayley J, Fossel AH, et al. Degenerative lumbar spinal stenosis. Diagnostic value of the history and physical examination. *Arthritis Rheum*. 1995;38:1236–41. [PMID: 7575718] [» CrossRef](#) [» Medline](#)


34.  Fayad F, Lefevre-Colau MM, Poiraudreau S, Fermanian J, Rannou F, Wlodyka Demaille S, et al. [Chronicity, recurrence, and return to work in low back pain: common prognostic factors]. *Ann Readapt Med Phys*. 2004;47:179–89. [PMID: 15130717] [» Medline](#)


35.  Pincus T, Burton AK, Vogel LI, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine*. 2002;27:E109–20. [PMID: 11880847] [» CrossRef](#) [» Medline](#)

36.  Gatchel RJ, Polatin PB, Noe C, Gardea M, Pulliam C, Thompson J. Treatment- and cost-effectiveness of early intervention for acute low-back pain patients: a one-year prospective study. *J Occup Rehabil*. 2003;13:1–9. [PMID: 12611026] [» CrossRef](#) [» Medline](#)


37.  Hay EM, Mullis R, Lewis M, Vohora K, Main CJ, Watson P, et al. Comparison of physical treatments versus a brief pain-management programme for back pain in primary care: a randomised clinical trial in physiotherapy practice. *Lancet*. 2005;365:2024–30. [PMID: 15950716] [» CrossRef](#) [» Medline](#)


38.  Jellema P, van der Windt DA, van der Horst HE, Twisk JW, Stalman WA, Bouter LM. Should treatment of (sub)acute low back pain be aimed at psychosocial prognostic factors? Cluster randomised clinical trial in general practice. *BMJ*. 2005;331:84. [PMID: 15967762] [» Abstract/FREE Full Text](#)

39.  Steenstra IA, Verbeek JH, Heymans MW, Bongers PM. Prognostic factors for duration of sick leave in patients sick listed with acute low back pain: a systematic review of the literature. *Occup Environ Med*. 2005;62:851–60. [PMID: 16299094] [» Abstract/FREE Full Text](#)

40.  Hestbaek L, Leboeuf-Yde C, Manniche C. Low back pain: what is the long-term course? A review of studies of general patient populations. *Eur Spine J*. 2003;12:149–65. [PMID: 12709853] [» Medline](#)

41.  Deyo RA, Diehl AK, Rosenthal M. Reducing roentgenography use. Can patient expectations be altered? *Arch Intern Med*. 1987;147:141–5. [PMID: 2948466] [» Abstract/FREE Full Text](#)

42.  Kendrick D, Fielding K, Bentley E, Kerslake R, Miller P, Pringle M. Radiography of the lumbar spine in primary care patients with low back pain: randomised controlled trial. *BMJ*. 2001;322:400–5. [PMID: 11179160] [» Abstract/FREE Full Text](#)

43.  Kerry S, Hilton S, Dundas D, Rink E, Oakeshott P. Radiography for low back pain: a randomised controlled trial and observational study in primary care. *Br J Gen Pract*. 2002;52:469–74. [PMID: 12051211] [» Medline](#)



VIEWED 03-16-2011

44. █ Jarvik JG. Imaging of adults with low back pain in the primary care setting. *Neuroimaging Clin N Am.* 2003;13:293–305. [PMID: 13677808]
» [CrossRef](#) » [Medline](#)

45. █ Gilbert F, Grant A, Gillan M, et al. Scottish Back Trial Group. Low back pain: influence of early MR imaging or CT on treatment and outcome—multicenter randomized trial. *Radiology.* 2004;231:343–51. [PMID: 15031430]

46. █ Jarvik JG, Hollingworth W, Martin B, Emerson SS, Gray DT, Overman S, et al. Rapid magnetic resonance imaging vs radiographs for patients with low back pain: a randomized controlled trial. *JAMA.* 2003;289:2810–8. [PMID: 12783911]
» [Abstract/FREE Full Text](#)

47. █ Lurie JD, Birkmeyer NJ, Weinstein JN. Rates of advanced spinal imaging and spine surgery. *Spine.* 2003;28:616–20. [PMID: 12642771]
» [CrossRef](#)
» [Medline](#)

48. █ Loblaw DA, Perry J, Chambers A, Laperriere NJ. Systematic review of the diagnosis and management of malignant extradural spinal cord compression: the Cancer Care Ontario Practice Guidelines Initiative's Neuro-Oncology Disease Site Group. *J Clin Oncol.* 2005;23:2028–37. [PMID: 15774794]
» [Abstract/FREE Full Text](#)

49. █ Todd NV. Cauda equina syndrome: the timing of surgery probably does influence outcome. *Br J Neurosurg.* 2005;19:301–6; discussion 307–8. [PMID: 16455534]

50. █ Tsiodras S, Falagas ME. Clinical assessment and medical treatment of spine infections. *Clin Orthop Relat Res.* 2006;444:38–50. [PMID: 16523126]
» [Medline](#)

51. █ Joines JD, McNutt RA, Carey TS, Deyo RA, Rouhani R. Finding cancer in primary care outpatients with low back pain: a comparison of diagnostic strategies. *J Gen Intern Med.* 2001;16:14–23. [PMID: 11251746]
» [CrossRef](#)
» [Medline](#)

52. █ Suarez-Almazor ME, Belseck E, Russell AS, Mackel JV. Use of lumbar radiographs for the early diagnosis of low back pain. Proposed guidelines would increase utilization. *JAMA.* 1997;277:1782–6. [PMID: 9178791]
» [Abstract/FREE Full Text](#)

53. █ Vroomen PC, de Krom MC, Knottnerus JA. Predicting the outcome of sciatica at short-term follow-up. *Br J Gen Pract.* 2002;52:119–23. [PMID: 11887877]
» [Medline](#)

54. █ Weber H. Lumbar disc herniation: A controlled prospective study with ten years of observation. *Spine.* 1983;8:131–40. [PMID: 6837383]
» [Medline](#)

55. █ Modic MT, Obuchowski NA, Ross JS, Brant-Zawadzki MN, Grooff PN, Mazanec DJ, et al. Acute low back pain and radiculopathy: MR imaging findings and their prognostic role and effect on outcome. *Radiology.* 2005;237:597–604. [PMID: 16244269]
» [Abstract/FREE Full Text](#)

56. █ Gibson JN, Grant IC, Waddell G. Surgery for lumbar disc prolapse. *Cochrane Database Syst Rev.* 2000;CD001350. [PMID: 10908492]
» [Medline](#)

57. █ Gibson JN, Waddell G. Surgery for degenerative lumbar spondylosis. *Cochrane Database Syst Rev.* 2005;CD001352. [PMID: 16235281]
» [Medline](#)

58. █ Nelemans PJ, deBie RA, deVet HC, Sturmans F. Injection therapy for subacute and chronic benign low back pain. *Spine.* 2001;26:501–15. [PMID: 11242378]
» [CrossRef](#) » [Medline](#)

59. █ Peul WC, van Houwelingen HC, van den Hout WB, et al. Leiden–The Hague Spine Intervention Prognostic Study Group. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med.* 2007;356:2245–56. [PMID: 17538084]
» [CrossRef](#) » [Medline](#)

60. █ Weinstein JN, Lurie JD, Tosteson TD, Skinner JS, Hanscom B, Tosteson AN, et al. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT) observational cohort. *JAMA.* 2006;296:2451–9. [PMID: 17119141]
» [Abstract/FREE Full Text](#)

61. █ Amundsen T, Weber H, Nordal HJ, Magnaes B, Abdelnoor M, Lilleås F. Lumbar spinal stenosis: conservative or surgical management?: A prospective 10-year study. *Spine.* 2000;25:1424–35; discussion 1435–6. [PMID: 10828926]

62. █ Atlas SJ, Keller RB, Wu YA, Deyo RA, Singer DE. Long-term outcomes of surgical and nonsurgical management of lumbar spinal stenosis: 8 to 10 year results from the Maine lumbar spine study. *Spine.* 2005;30:936–43. [PMID: 15834339]
» [CrossRef](#) » [Medline](#)

63. █ Weinstein JN, Lurie JD, Tosteson TD, Hanscom B, Tosteson AN, Blood EA, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med.* 2007;356:2257–70. [PMID: 17538085]
» [CrossRef](#) » [Medline](#)

64. █ Malmivaara A, Slati P, Heliovaara M, et al. Finnish Lumbar Spinal Research Group. Surgical or nonoperative treatment for lumbar spinal stenosis?

VIEWED 03-16-2011

A randomized controlled trial. *Spine*. 2007;32:1–8. [PMID: 17202885]
» [CrossRef](#) » [Medline](#)

65. Hagen KB, Hilde G, Jamtvedt G, Winnem M. Bed rest for acute low-back pain and sciatica. *Cochrane Database Syst Rev*. 2004;CD001254. [PMID: 15495012] » [Medline](#)

66. Hilde G, Hagen KB, Jamtvedt G, Winnem M. Advice to stay active as a single treatment for low back pain and sciatica. *Cochrane Database Syst Rev*. 2002;CD003632. [PMID: 12076492] » [Medline](#)

67. Burton AK, Waddell G, Tillotson KM, Summerton N. Information and advice to patients with back pain can have a positive effect. A randomized controlled trial of a novel educational booklet in primary care. *Spine*. 1999;24:2484–91. [PMID: 10626311] » [CrossRef](#) » [Medline](#)

68. Cherkin DC, Deyo RA, Battié M, Street J, Barlow W. A comparison of physical therapy, chiropractic manipulation, and provision of an educational booklet for the treatment of patients with low back pain. *N Engl J Med*. 1998;339:1021–9. [PMID: 9761803] » [CrossRef](#) » [Medline](#)

69. Cherkin DC, Eisenberg D, Sherman KJ, Barlow W, Kaptchuk TJ, Street J, et al. Randomized trial comparing traditional Chinese medical acupuncture, therapeutic massage, and self-care education for chronic low back pain. *Arch Intern Med*. 2001;161:1081–8. [PMID: 11322842] » [Abstract/FREE Full Text](#)

70. Sherman KJ, Cherkin DC, Erro J, Miglioretti DL, Deyo RA. Comparing yoga, exercise, and a self-care book for chronic low back pain: a randomized, controlled trial. *Ann Intern Med*. 2005;143:849–56. [PMID: 16365466]
» [Abstract/FREE Full Text](#)

71. Scheel IB, Hagen KB, Herrin J, Carling C, Oxman AD. Blind faith? The effects of promoting active sick leave for back pain patients: a cluster-randomized controlled trial. *Spine*. 2002;27:2734–40. [PMID: 12461401]
» [CrossRef](#) » [Medline](#)

72. Indahl A, Velund L, Reikeraas O. Good prognosis for low back pain when left untampered. A randomized clinical trial. *Spine*. 1995;20:473–7. [PMID: 7747232] » [Medline](#)

73. Karjalainen K, Malmivaara A, Pohjolainen T, Hurri H, Mutanen P, Rissanen P, et al. Mini-intervention for subacute low back pain: a randomized controlled trial. *Spine*. 2003;28:533–40; discussion 540–1. [PMID: 12642757]

74. Hagen EM, Eriksen HR, Ursin H. Does early intervention with a light mobilization program reduce long-term sick leave for low back pain? *Spine*. 2000;25:1973–6. [PMID: 10908942] » [CrossRef](#) » [Medline](#)

75. French SD, Cameron M, Walker BF, Reggars JW, Esterman AJ. Superficial heat or cold for low back pain. *Cochrane Database Syst Rev*. 2006;CD004750. [PMID: 16437495] » [Medline](#)

76. Kovacs FM, Abaira V, Peña A, Martín-Rodríguez JG, Sánchez-Vera M, Ferrer E, et al. Effect of firmness of mattress on chronic non-specific low-back pain: randomised, double-blind, controlled, multicentre trial. *Lancet*. 2003;362:1599–604. [PMID: 14630439] » [CrossRef](#) » [Medline](#)

77. Jellema P, van Tulder MW, van Poppel MN, Nachemson AL, Bouter LM. Lumbar supports for prevention and treatment of low back pain: a systematic review within the framework of the Cochrane Back Review Group. *Spine*. 2001;26:377–86. [PMID: 11224885] » [CrossRef](#) » [Medline](#)

78. Lee C, Straus WL, Balshaw R, Barlas S, Vogel S, Schnitzer TJ. A comparison of the efficacy and safety of nonsteroidal antiinflammatory agents versus acetaminophen in the treatment of osteoarthritis: a meta-analysis. *Arthritis Rheum*. 2004;51:746–54. [PMID: 15478167] » [CrossRef](#) » [Medline](#)

79. Towheed TE, Judd MJ, Hochberg MC, Wells G. Acetaminophen for osteoarthritis. *Cochrane Database Syst Rev*. 2003;CD004257. [PMID: 12804508] » [Medline](#)

80. van Tulder MW, Scholten RJ, Koes BW, Deyo RA. Nonsteroidal anti-inflammatory drugs for low back pain: a systematic review within the framework of the Cochrane Collaboration Back Review Group. *Spine*. 2000;25:2501–13. [PMID: 11013503] » [CrossRef](#) » [Medline](#)

81. Wegman A, van der Windt D, van Tulder M, Stalman W, de Vries T. Nonsteroidal antiinflammatory drugs or acetaminophen for osteoarthritis of the hip or knee? A systematic review of evidence and guidelines. *J Rheumatol*. 2004;31:344–54. [PMID: 14760807] » [Abstract/FREE Full Text](#)

82. Zhang W, Jones A, Doherty M. Does paracetamol (acetaminophen) reduce the pain of osteoarthritis? A meta-analysis of randomised controlled trials. *Ann Rheum Dis*. 2004;63:901–7. [PMID: 15020311] » [Abstract/FREE Full Text](#)

83. Hernández-Díaz S, Rodríguez LA. Association between nonsteroidal anti-inflammatory drugs and upper gastrointestinal tract bleeding/perforation: an overview of epidemiologic studies published in the 1990s. *Arch Intern Med*. » [Abstract/FREE Full Text](#)

VIEWED 03-16-2011

2000;160:2093–9. [PMID: 10904451]

84. Rahme E, Pettitt D, LeLorier J. Determinants and sequelae associated with utilization of acetaminophen versus traditional nonsteroidal antiinflammatory drugs in an elderly population. *Arthritis Rheum.* 2002;46:3046–54. [PMID: 12428249] »CrossRef »Medline

85. Watkins PB, Kaplowitz N, Slattery JT, Colonese CR, Colucci SV, Stewart PW, et al. Aminotransferase elevations in healthy adults receiving 4 grams of acetaminophen daily: a randomized controlled trial. *JAMA.* 2006;296:87–93. [PMID: 16820551] »Abstract/FREE Full Text

86. Kearney PM, Baigent C, Godwin J, Halls H, Emberson JR, Patrono C. Do selective cyclo-oxygenase-2 inhibitors and traditional non-steroidal anti-inflammatory drugs increase the risk of atherothrombosis? Meta-analysis of randomised trials. *BMJ.* 2006;332:1302–8. [PMID: 16740558] »Abstract/FREE Full Text

87. Lai KC, Chu KM, Hui WM, Wong BC, Hu WH, Wong WM, et al. Celecoxib compared with lansoprazole and naproxen to prevent gastrointestinal ulcer complications. *Am J Med.* 2005;118:1271–8. [PMID: 16271912] »CrossRef »Medline

88. Derry S, Loke YK. Risk of gastrointestinal haemorrhage with long term use of aspirin: meta-analysis. *BMJ.* 2000;321:1183–7. [PMID: 11073508] »Abstract/FREE Full Text

89. Furlan AD, Sandoval JA, Mailis-Gagnon A, Tunks E. Opioids for chronic noncancer pain: a meta-analysis of effectiveness and side effects. *CMAJ.* 2006;174:1589–94. [PMID: 16717269] »Abstract/FREE Full Text

90. Kalso E, Edwards JE, Moore RA, McQuay HJ. Opioids in chronic non-cancer pain: systematic review of efficacy and safety. *Pain.* 2004;112:372–80. [PMID: 15561393] »CrossRef »Medline

91. Martell BA, O'Connor PG, Kerns RD, Becker WC, Morales KH, Kosten TR, et al. Systematic review: opioid treatment for chronic back pain: prevalence, efficacy, and association with addiction. *Ann Intern Med.* 2007;146:116–27. [PMID: 17227935] »Abstract/FREE Full Text

92. Collins A, Simpson K, eds. Recommendations for the Appropriate Use of Opioids for Persistent Non-Cancer Pain. London: The Pain Society; 2005.

93. Jovey R, Ennis J, Garder-Nix J, Goldman B, Hayes H, Lynch M, et al.; Canadian Pain Society. Use of opioid analgesics for the treatment of chronic noncancer pain—a consensus statement and guidelines from the Canadian Pain Society, 2002. *Pain Res Manag.* 2005;8 Suppl A:3A–28A. [PMID: 14685504] »Medline

94. Kalso E, Allan L, DelleMijn PL, Faura CC, Ilias WK, Jensen TS, et al. Recommendations for using opioids in chronic non-cancer pain. *Eur J Pain.* 2003;7:381–6. [PMID: 12935789] »CrossRef »Medline

95. Chou R, Clark E, Helfand M. Comparative efficacy and safety of long-acting oral opioids for chronic non-cancer pain: a systematic review. *J Pain Symptom Manage.* 2003;26:1026–48. [PMID: 14585554] »CrossRef »Medline

96. van Tulder M, Touray T, Furlan A, Solway S, Bouter L. Cochrane Back Review Group. Muscle relaxants for nonspecific low back pain: a systematic review within the framework of the Cochrane Collaboration. *Spine.* 2003;28:1978–92. [PMID: 12973146] »CrossRef »Medline

97. Chou R, Peterson K, Helfand M. Comparative efficacy and safety of skeletal muscle relaxants for spasticity and musculoskeletal conditions: a systematic review. *J Pain Symptom Manage.* 2004;28:140–75. [PMID: 15276195] »CrossRef »Medline

98. Salerno SM, Browning R, Jackson JL. The effect of antidepressant treatment on chronic back pain: a meta-analysis. *Arch Intern Med.* 2002;162:19–24. [PMID: 11784215] »Abstract/FREE Full Text

99. Staiger TO, Gaster B, Sullivan MD, Deyo RA. Systematic review of antidepressants in the treatment of chronic low back pain. *Spine.* 2003;28:2540–5. [PMID: 14624092] »CrossRef »Medline

100. Bair MJ, Robinson RL, Katon W, Kroenke K. Depression and pain comorbidity: a literature review. *Arch Intern Med.* 2003;163:2433–45. [PMID: 14609780] »Abstract/FREE Full Text


101. McClean G. Does gabapentin have an analgesic effect on background, movement and referred pain? A randomised, double-blind, placebo controlled study. *The Pain Clinic.* 2001;13:103–7.


102. Yildirim K, Sisecioglu M, Karatay S, et al. The effectiveness of gabapentin in patients with chronic radiculopathy. *The Pain Clinic.* 2003;15:213–8.


103. Gagnier JJ, van Tulder M, Berman B, Bombardier C. Herbal medicine for


VIEWED 03-16-2011


low back pain. Cochrane Database Syst Rev. 2006;CD004504. [PMID: 16625605] [» Medline](#)


104.  Finckh A, Zufferey P, Schurch MA, Balagué F, Waldburger M, So AK. Short-term efficacy of intravenous pulse glucocorticoids in acute discogenic sciatica. A randomized controlled trial. Spine. 2006;31:377–81. [PMID: 16481946] [» Medline](#)


105.  Friedman BW, Holden L, Esses D, Bijur PE, Choi HK, Solorzano C, et al. Parenteral corticosteroids for Emergency Department patients with non-radicular low back pain. J Emerg Med. 2006;31:365–70. [PMID: 17046475] [» CrossRef](#) [» Medline](#)


106.  Haimovic IC, Beresford HR. Dexamethasone is not superior to placebo for treating lumbosacral radicular pain. Neurology. 1986;36:1593–4. [PMID: 2946981] [» Abstract/FREE Full Text](#)


107.  Porsman O, Friis H. Prolapsed lumbar disc treated with intramuscularly administered dexamethasonephosphate. A prospectively planned, double-blind, controlled clinical trial in 52 patients. Scand J Rheumatol. 1979;8:142–4. [PMID: 386492] [» Medline](#)


108.  Assendelft WJ, Morton SC, Yu EI, Suttrop MJ, Shekelle PG. Spinal manipulative therapy for low back pain. A meta-analysis of effectiveness relative to other therapies. Ann Intern Med. 2003;138:871–81. [PMID: 12779297] [» Abstract/FREE Full Text](#)


109.  Hayden JA, van Tulder MW, Tomlinson G. Systematic review: strategies for using exercise therapy to improve outcomes in chronic low back pain. Ann Intern Med. 2005;142:776–85. [PMID: 15867410] [» Abstract/FREE Full Text](#)


110.  Waddell G, McIntosh A, Hutchinson A, Feder G, Lewis M. Clinical Guidelines for the Management of Acute Low Back Pain: Low Back Pain Evidence Review. London: Royal College of General Practitioners; 1996.


111.  Karjalainen K, Malmivaara A, van Tulder M, Roine R, Jauhiainen M, Hurri H, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain in working-age adults: a systematic review within the framework of the Cochrane Collaboration Back Review Group. Spine. 2001;26:262–9. [PMID: 11224862] [» CrossRef](#) [» Medline](#)


112.  Schonstein E, Kenny D, Keating J, Koes B, Herbert RD. Physical conditioning programs for workers with back and neck pain: a cochrane systematic review. Spine. 2003;28:E391–5. [PMID: 14520951] [» Medline](#)


113.  Pengel HM, Maher CG, Refshauge KM. Systematic review of conservative interventions for subacute low back pain. Clin Rehabil. 2002;16:811–20. [PMID: 12501942] [» Abstract/FREE Full Text](#)


114.  Furlan AD, van Tulder M, Cherkin D, Tsukayama H, Lao L, Koes B, et al. Acupuncture and dry-needling for low back pain: an updated systematic review within the framework of the cochrane collaboration. Spine. 2005;30:944–63. [PMID: 15834340] [» CrossRef](#) [» Medline](#)


115.  Manheimer E, White A, Berman B, Forsy K, Ernst E. Meta-analysis: acupuncture for low back pain. Ann Intern Med. 2005;142:651–63. [PMID: 15838072] [» Abstract/FREE Full Text](#)


116.  Furlan AD, Brosseau L, Imamura M, Irvin E. Massage for low-back pain: a systematic review within the framework of the Cochrane Collaboration Back Review Group. Spine. 2002;27:1896–910. [PMID: 12221356] [» CrossRef](#) [» Medline](#)


117.  Hoffman BM, Papas RK, Chatkoff DK, Kerns RD. Meta-analysis of psychological interventions for chronic low back pain. Health Psychol. 2007;26:1–9. [PMID: 17209691] [» CrossRef](#) [» Medline](#)


118.  Ostelo RW, van Tulder MW, Vlaeyen JW, Linton SJ, Morley SJ, Assendelft WJ. Behavioural treatment for chronic low-back pain. Cochrane Database Syst Rev. 2005;CD002014. [PMID: 15674889] [» Medline](#)

119.  Guzmán J, Esmail R, Karjalainen K, Malmivaara A, Irvin E, Bombardier C. Multidisciplinary rehabilitation for chronic low back pain: systematic review. BMJ. 2001;322:1511–6. [PMID: 11420271] [» Abstract/FREE Full Text](#)

120.  Clarke J, van Tulder M, Blomberg S, de Vet H, van der Heijden G, Bronfort G. Traction for low back pain with or without sciatica: an updated systematic review within the framework of the Cochrane collaboration. Spine. 2006;31:1591–9. [PMID: 16778694] [» Medline](#)

121.  Harte AA, Baxter GD, Gracey JH. The efficacy of traction for back pain: a systematic review of randomized controlled trials. Arch Phys Med Rehabil. 2003;84:1542–53. [PMID: 14586924] [» CrossRef](#) [» Medline](#)

122.  Vroomen PC, de Krom MC, Slofstra PD, Knottnerus JA. Conservative treatment of sciatica: a systematic review. J Spinal Disord. 2000;13:463–9. [PMID: 11132976] [» CrossRef](#) [» Medline](#)

123.  Kalauokalani D, Cherkin DC, Sherman KJ, Koepsell TD, Deyo RA.

VIEWED 03-16-2011

Lessons from a trial of acupuncture and massage for low back pain: patient expectations and treatment effects. Spine. 2001;26:1418–24. [PMID: 11458142] [»CrossRef](#) [»Medline](#)

124. Childs JD, Fritz JM, Flynn TW, Irrgang JJ, Johnson KK, Majkowski GR, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. Ann Intern Med. 2004;141:920–8. [PMID: 15611489] [»Abstract/FREE Full Text](#)

125. Brennan GP, Fritz JM, Hunter SJ, Thackeray A, Delitto A, Erhard RE. Identifying subgroups of patients with acute/subacute “nonspecific” low back pain: results of a randomized clinical trial. Spine. 2006;31:623–31. [PMID: 16540864] [»CrossRef](#) [»Medline](#)

126. Fritz JM, Delitto A, Erhard RE. Comparison of classification-based physical therapy with therapy based on clinical practice guidelines for patients with acute low back pain: a randomized clinical trial. Spine. 2003;28:1363–71; discussion 1372. [PMID: 12838091]

127. Heymans MW, van Tulder MW, Esmail R, Bombardier C, Koes BW. Back schools for nonspecific low back pain: a systematic review within the framework of the Cochrane Collaboration Back Review Group. Spine. 2005;30:2153–63. [PMID: 16205340] [»CrossRef](#) [»Medline](#)

128. Airaksinen O, Brox J, Cedraschi C, et al. COST B13 Working Group on Guidelines for Chronic Low Back Pain. Chapter 4. European guidelines for the management of chronic nonspecific low back pain. Eur Spine J. 2006;15 Suppl 2:S192–300. [PMID: 16550448] [»CrossRef](#) [»Medline](#)

129. Brox JI, Sørensen R, Friis A, Nygaard Ø, Indahl A, Keller A, et al. Randomized clinical trial of lumbar instrumented fusion and cognitive intervention and exercises in patients with chronic low back pain and disc degeneration. Spine. 2003;28:1913–21. [PMID: 12973134] [»CrossRef](#) [»Medline](#)

130. Fairbank J, Frost H, Wilson–MacDonald J, Yu LM, Barker K, Collins R; Spine Stabilisation Trial Group. Randomised controlled trial to compare surgical stabilisation of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilisation trial. BMJ. 2005;330:1233. [PMID: 15911537] [»Abstract/FREE Full Text](#)


131. Fritzell P, Hagg O, Wessberg P, Nordwall A. Swedish Lumbar Spine Study Group. 2001 Volvo Award Winner in Clinical Studies: Lumbar fusion versus nonsurgical treatment for chronic low back pain: a multicenter randomized controlled trial from the Swedish Lumbar Spine Study Group. Spine. 2001;26:2521–32; discussion 2532–4. [PMID: 11725230]

VIEWED 03-16-2011

 CiteULike

 Complere

 Connotea


 Del.icio.us

 Digg

 Facebook

 Reddit

 Technorati

 Twitter

What's this?

Related articles

Clinical Guidelines:
Nonpharmacologic Therapies for Acute and Chronic Low Back Pain: A Review of the Evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline
Roger Chou and Laurie Hoyt Huffman
Ann Intern Med October 2, 2007 147:492–504
[»Abstract](#) [»Full Text](#) [»Full Text \(PDF\)](#) [»Video News Release](#)

Clinical Guidelines:
Medications for Acute and Chronic Low Back Pain: A Review of the Evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline
Roger Chou and Laurie Hoyt Huffman
Ann Intern Med October 2, 2007 147:505–514
[»Abstract](#) [»Full Text](#) [»Full Text \(PDF\)](#) [»Video News Release](#)

Letter:
Overviews and Systematic Reviews on Low Back Pain
Jan M. Bjordal, Atle Klovning, Rodrigo Alvaro B. Lopes–Martins, Pål–Didrik Hoff Roland, Jon Joensen, and Lars Slørdal
Ann Intern Med May 20, 2008 148:789–790
[»Excerpt](#) [»Full Text](#) [»Full Text \(PDF\)](#)

Letter:
Overviews and Systematic Reviews on Low Back Pain
Edzard Ernst
Ann Intern Med May 20, 2008 148:790–791
[»Excerpt](#) [»Full Text](#) [»Full Text \(PDF\)](#)

Letter:
Overviews and Systematic Reviews on Low Back Pain
Roger Chou, Paul Shekelle, Amir Qaseem, and Douglas K. Owens
Ann Intern Med May 20, 2008 148:791–792
[»Excerpt](#) [»Full Text](#) [»Full Text \(PDF\)](#)

Comments on This Article

Re: Early Physical Therapy intervention necessary for low back pain
Amir Qaseem and Roger Chou, MD, Douglas K. Owens, MD, MS
Ann Intern Med published online August 28, 2008
[»Full Text](#)

Early Physical Therapy intervention necessary for low back pain.
Philip P Tygiel and
Britt Smith PT, DPT, OCS, FAAOMPT, Eric Robertson PT, DPT, Mark Shropshire PT
Ann Intern Med published online May 12, 2008
[»Full Text](#)

Middle cluneal neuropathy: a neglected cause of low back pain.
Stanley J Antolak
Ann Intern Med published online December 5, 2007
[»Full Text](#)

Correction
Roger Chou and
Roger Chou, Amir Qaseem, Vincenza Snow and Douglas Owens
Ann Intern Med published online December 5, 2007
[»Full Text](#)

In Response
Roger Chou and
Roger Chou, Amir Qaseem, Vincenza Snow and Douglas Owens
Ann Intern Med published online December 5, 2007
[»Full Text](#)

Failing evidence behind first-line medication recommendations
Jan M Bjordal and
Atle Klovning, Rodrigo Lopes-Martins, Lars Slørdal, Jon Joensen, Pål-Didrik H. R
Ann Intern Med published online November 7, 2007
[»Full Text](#)

Summary for Patients

Summaries for Patients:
Diagnosis and Treatment of Low Back Pain: Recommendations from
the American College of Physicians/American Pain Society
Ann Intern Med October 2, 2007 147:1–45
[»Excerpt](#) [»Full Text](#) [»Full Text \(PDF\)](#) [»Video News Release](#)

Articles citing this article

Depressive Symptoms, Anatomical Region, and Clinical Outcomes for
Patients Seeking Outpatient Physical Therapy for Musculoskeletal Pain
ptjournal March 1, 2011 91:358–372
[»Abstract](#) [»Full Text](#) [»Full Text \(PDF\)](#)

VIEWED 03-16-2011

High-Value, Cost-Conscious Health Care: Concepts for Clinicians to Evaluate the Benefits, Harms, and Costs of Medical Interventions
ANN INTERN MED February 1, 2011 154:174-180
»Abstract »Full Text »Full Text (PDF)

Diagnostic Imaging for Low Back Pain: Advice for High-Value Health Care From the American College of Physicians
ANN INTERN MED February 1, 2011 154:181-189
»Abstract »Full Text »Full Text (PDF)

Engaging Physicians in Change: Results of a Safety Net Quality Improvement Program to Reduce Overuse
American Journal of Medical Quality January 1, 2011 26:26-33
»Abstract »Full Text (PDF)

Pain Management and the Primary Care Encounter: Opportunities For Quality Improvement
Journal of Primary Care & Community Health January 1, 2011 2:37-44
»Abstract »Full Text (PDF)

The diagnosis and management of common non-specific back pain – a clinical review
Trauma January 1, 2011 13:57-64
»Abstract »Full Text (PDF)

Does This Older Adult With Lower Extremity Pain Have the Clinical Syndrome of Lumbar Spinal Stenosis?
JAMA December 15, 2010 304:2628-2636
»Abstract »Full Text »Full Text (PDF)

How the AOA Established the First National Guidelines for OMT
J Am Osteopath Assoc November 1, 2010 110:36-37
»Full Text »Full Text (PDF)

Patients' Knowledge about 9 Common Health Conditions: The DECISIONS Survey
Med Decis Making September 1, 2010 30:355-525
»Abstract »Full Text (PDF)

Lumbar Spine MRI for Low Back Pain: Indications and Yield
Am. J. Roentgenol. September 1, 2010 195:550-559
»Abstract »Full Text »Full Text (PDF)

A guide to imaging for common neurological problems
BMJ August 16, 2010 341:c4113
»Full Text

Article Commentary: Response to Commentary on Clinical practice guideline: Hoarseness (Dysphonia)
Otolaryngology -- Head and Neck Surgery August 1, 2010 143:178-180
»Abstract »Full Text »Full Text (PDF)

Glucosamine and the Ongoing Enigma of Chronic Low Back Pain
JAMA July 7, 2010 304:93-94
»Full Text »Full Text (PDF)

Clinical significance of inflammatory back pain for diagnosis and screening of patients with axial spondyloarthritis
Ann Rheum Dis July 1, 2010 69:1264-1268
»Abstract »Full Text »Full Text (PDF)

Magnetic resonance imaging for low back pain: indications and

VIEWED 03-16-2011

limitations

Postgrad. Med. J. June 1, 2010 86:374–378

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Perceived Benefit of Complementary and Alternative Medicine (CAM) for Back Pain: A National Survey

J Am Board Fam Med May 1, 2010 23:354–362

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Spondyloarthritis in a Patient With Unilateral Buttock Pain and History of Crohn Disease

ptjournal May 1, 2010 90:784–792

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Will This Patient Develop Persistent Disabling Low Back Pain?

JAMA April 7, 2010 303:1295–1302

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Pursuit and Implementation of Hospital–Based Outpatient Direct Access to Physical Therapy Services: An Administrative Case Report

ptjournal January 1, 2010 90:100–109

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Magnetic resonance imaging for low back pain: indications and limitations

Ann Rheum Dis January 1, 2010 69:7–11

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Simulated acupuncture and real acupuncture both reduced low back pain dysfunction and symptoms

Evid. Based Med. December 1, 2009 14:179

»[Full Text](#) »[Full Text \(PDF\)](#)

Motor Control Exercise for Chronic Low Back Pain: A Randomized Placebo–Controlled Trial

ptjournal December 1, 2009 89:1275–1286

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Magnetic Resonance Imaging And Low Back Pain Care For Medicare Patients

Health Aff (Millwood) November 1, 2009 28:w1133–w1140

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

An Evidence–Based Approach to Spine Surgery

American Journal of Medical Quality November 1, 2009 24:155–245

»[Abstract](#) »[Full Text \(PDF\)](#)

A Comparison of Long– and Short–Acting Opioids for the Treatment of Chronic Noncancer Pain: Tailoring Therapy to Meet Patient Needs

Mayo Clin Proc. July 1, 2009 84:602–612

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Back pain made simple: An approach based on principles and evidence

Cleveland Clinic Journal of Medicine July 1, 2009 76:393–399

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Rapidity and Modality of Imaging for Acute Low Back Pain in Elderly Patients

Arch Intern Med May 25, 2009 169:972–981

»[Abstract](#) »[Full Text](#) »[Full Text \(PDF\)](#)

Imaging Idolatry: The Uneasy Intersection of Patient Satisfaction, Quality of Care, and Overuse

VIEWED 03-16-2011

Arch Intern Med May 25, 2009 169:921–923

»Full Text »Full Text (PDF)

Analgesic effects of treatments for non-specific low back pain: a meta-analysis of placebo-controlled randomized trials
Rheumatology (Oxford) May 1, 2009 48:520–527

»Abstract »Full Text »Full Text (PDF)

Small effects of treatments for non-specific low back pain: how can we improve patients' outcomes?

Rheumatology (Oxford) May 1, 2009 48:464–465

»Full Text »Full Text (PDF)

Overtreating Chronic Back Pain: Time to Back Off?

J Am Board Fam Med January 1, 2009 22:62–68

»Abstract »Full Text »Full Text (PDF)

Bibliography: Geriatric Psychiatry

Focus January 1, 2009 7:46–48

»Full Text »Full Text (PDF)

Management of low back pain

BMJ December 22, 2008 337:a2718

»Full Text

Prescription Opioids, Overdose Deaths, and Physician Responsibility

JAMA December 10, 2008 300:2672–2673

»Full Text »Full Text (PDF)

Overviews and Systematic Reviews on Low Back Pain

ANN INTERN MED May 20, 2008 148:799–799

»Full Text »Full Text (PDF)

Overviews and Systematic Reviews on Low Back Pain

ANN INTERN MED May 20, 2008 148:791–792

»Full Text »Full Text (PDF)

A 50-Year-Old Man With Chronic Low Back Pain

JAMA May 7, 2008 299:2066–2077

»Abstract »Full Text »Full Text (PDF)

Correction: Diagnosis and Treatment of Low Back Pain

ANN INTERN MED February 5, 2008 148:247–248

»Full Text »Full Text (PDF)

Diagnosis and Treatment of Low Back Pain

JWatch General October 25, 2007 2007:1

»Full Text

VIEWED 03-16-2011

VIEWED 03-16-2011

VIEWED 03-16-2011